

The Fadin-Lipatov-vertex in high-energy heavy quark and charmonium production[★]

Philipp Hägler^a

^a*Universität Regensburg, Institut für theoretische Physik,
93040 Regensburg, Germany*

Abstract

Calculations of heavy quark and charmonium hadroproduction are presented in the $k_{\perp}$ -factorization approach. The resulting differential cross sections are compared to experimental values and NLO collinear predictions. The application of the full effective Fadin-Lipatov- $q\bar{q}$ -production-vertex has an important influence on the results and in particular leads to a strong suppression of certain color octet contributions which are related to the outstanding problem of the production of polarized J/Ψ .

1 Heavy quark production

At very high energies $\sqrt{s} \rightarrow \infty$ in the small- x -region the Multi-Regge-Kinematics (MRK) respectively the Quasi-MRK (QMRK) is an especially suited approximation for the calculation of lepto- and hadroproduction cross sections. We consider the central heavy quark-antiquark-hadroproduction process in figs. 1,2 and choose the frame where the proton momenta are given by

$$P_1^+ \equiv P_1^0 + P_1^3 = P_2^- = \sqrt{s}, \quad P_1^- = P_2^+ = P_{1,2}^{\perp} = 0.$$

In the QMRK we have then an almost transverse momentum transfer $t_{1,2} \approx q_{1,2\perp}^2$ and for the t-channel momenta the relations

$$q_1^+ \gg q_2^+, \quad q_2^- \gg q_1^-,$$

but in general no strong ordering of the momenta of the produced heavy quark and antiquark $k_1 \approx k_2$. This kinematics corresponds to the production

[★] This talk is based on results obtained in collaboration with R. Kirschner, A. Schäfer, L. Szymanowski and O.V. Teryaev.




Fig. 1. Production of a $q\bar{q}$ -cluster in QMRK

of a $q\bar{q}$ -cluster with finite invariant mass and therefore makes the production of a $q\bar{q}$ -bound state possible. The associated factorization scheme is the $k_\perp$ -factorization and the particles in the t-channel are allowed to have finite transverse momenta. The cross section for the inclusive hadroproduction of a $q\bar{q}$ -pair at very high energies where gluon exchange dominates is then given by [1,2]

$$\sigma_{P_1 P_2 \rightarrow q\bar{q} X} = \frac{1}{16 (2\pi)^4} \int \frac{d^3 k_1}{k_1^+} \frac{d^3 k_2}{k_2^+} d^2 q_{1\perp} d^2 q_{2\perp} \delta^2 (q_{1\perp} - q_{2\perp} - k_{1\perp} - k_{2\perp}) \times \frac{\mathcal{F}(x_1, q_{1\perp})}{(q_{1\perp}^2)^2} \frac{(\psi^{c_2 c_1})^\dagger \psi^{c_2 c_1}}{(N_c^2 - 1)^2} \frac{\mathcal{F}(x_2, q_{2\perp})}{(q_{2\perp}^2)^2}. \quad (1)$$

The most important parts of this equation are the hard scattering amplitude




Fig. 2. $q\bar{q}$ -production in $k_\perp$ -factorization

$\psi^{c_2 c_1}$ and the unintegrated gluon distribution function $\mathcal{F}(x_{1,2}, q_{1,2\perp})$, where the longitudinal momentum fractions are defined as usual, $x_1 \equiv q_1^+/P_1^+$ and $x_2 \equiv -q_2^-/P_2^-$. Due to the QMRK the gluons are off-shell and longitudinal, which corresponds to the replacement $g^{\mu\nu} \approx 1/(2s) P_2^\mu P_1^\nu$ e.g. for the numerator of the upper gluon propagator. The use of the QMRK and off-shell gluons together with the requirement of gauge invariance determines the $q\bar{q}$ -production amplitude $\psi^{c_2 c_1}$ in terms of the NLLA BFKL production vertex $\Psi^{c_2 c_1}$, which is explicitly given by Fadin and Lipatov in [3]

$$\Psi^{c_2 c_1} \equiv \Psi^{c_2 c_1 - +} = -g^2 \left(T^{c_1} T^{c_2} b^{-+}(k_1, k_2) - T^{c_2} T^{c_1} b^{-+T}(k_2, k_1) \right),$$




Fig. 3. Bottom quark production

where the function $b^{-+}(k_1, k_2)$ is defined through

$$b^{-+}(k_1, k_2) \equiv \gamma^- \frac{\not{k}_{1\perp} - \not{k}_{1\perp} - m}{(q_1 - k_1)^2 - m^2} \gamma^+ - \frac{\gamma_\beta \Gamma^{+-\beta}(q_2, q_1)}{(k_1 + k_2)^2}, \quad (2)$$

and a very similar equation holds for the crossed contribution $b^{-+T}(k_2, k_1)$. Equation (2) includes in the second part on the right hand side the effective gluon-production-vertex

$$\Gamma^{+-\beta}(q_2, q_1) = \gamma^{+-\beta}(q_1, q_2) + \left\{ -2t_1 \frac{n^{-\beta}}{q_1^- - q_2^-} + 2t_2 \frac{n^{+\beta}}{q_1^+ - q_2^+} \right\}. \quad (3)$$

The first term of the right hand side of this equation is the standard three-gluon-coupling, while the second term in curly brackets is an additional part which is essential for gauge invariance. The function $\Gamma^{+-\beta}(q_2, q_1)$ satisfies the Ward-Slavnov-identity $(q_1 - q_2)_\beta \Gamma^{+-\beta}(q_2, q_1) = 0$, and gauge invariance leads to the vanishing of the amplitude $\psi^{c_2 c_1} = \bar{u}(k_1) \Psi^{c_2 c_1 -+} v(k_2) \xrightarrow{q_{1/2\perp} \rightarrow 0} 0$ in the limit of vanishing transverse momentum.

In order to be able to perform a reasonable comparison to experimental data we use the unintegrated gluon distribution function $\mathcal{F}(x, q_\perp)$ which has been calculated in [4]. The authors extracted $\mathcal{F}(x, q_\perp)$ from a fit to F_2 data and obtained an excellent χ^2/NDOF with the use of the following integrated gluon distribution at an input scale of $Q_0^2 = 1 \text{ GeV}^2$

$$xg(x, Q_0^2 = 1 \text{ GeV}^2) = 1.57(1 - x)^{2.5} \quad (4)$$




Fig. 4. $b\bar{b}$ -correlations

which enters the calculation through the relation

$$g(x, Q^2) = \left(\int_0^{Q_0^2} + \int_{Q_0^2}^{Q^2} \right) dk_{\perp}^2 \frac{\mathcal{F}(x, k_{\perp}^2)}{k_{\perp}^2} = xg(x, Q_0^2) + \int_{Q_0^2}^{Q^2} dk_{\perp}^2 \frac{\mathcal{F}(x, k_{\perp}^2)}{k_{\perp}^2}. \quad (5)$$

For a correct treatment of the region of small transverse gluon momenta $|q_{\perp}^2| < Q_0^2$ we have to modify the cross section formular (1) in a way which is compatible with equation (5) and the vanishing of the amplitude in the limit $q_{1/2\perp} \rightarrow 0$, for details see [2]. Our result for the production of a single bottom quark is shown in fig. 3 in comparison to data from the Tevatron collider [5] and NLO calculations in the collinear approach. The error band is due to a simultaneous variation of the bottom quark mass $4.5 \text{ GeV} < m_b < 4.9 \text{ GeV}$ and the fundamental scale $100 \text{ MeV} < \lambda_{QCD}^{n_f=5} < 180 \text{ MeV}$, while the scale dependence of the coupling constant $\alpha_S(\mu^2)$ is chosen to be $\mu^2 = |q_{1,2\perp}^2| + m^2$. We find in contrast to the NLO collinear calculation a good agreement between the $k_{\perp}$ -factorization prediction and the experimental values over the whole range of $k_{1\perp}$. Likewise important is the comparison to measurements of $b\bar{b}$ -correlations [6] in fig. 4. Although the cross section (1) is only of LO in α_S , the $k_{\perp}$ -factorization together with the effective production vertex and off-shell gluons gives a very good desription of the data which lie about a factor of 3 above the NLO collinear result.

2 Charmonium production

Motivated by the results for heavy quark production we now pass over to the hadroproduction of charmonia which is a far more controversial subject [7,8]. We consider the production of the χ_c , which is a 3P_J -state and the J/Ψ ,




Fig. 5. χ_c -meson-production

which is a 3S_1 state in the framework of the non-relativistic-QCD (NRQCD) approach and have therefore to include the color singlet and color octet contributions to the final color singlet bound states. The calculation is again based on the cross section formula (1) and the unintegrated gluon distribution from [4]. The details of the calculations and the binding mechanisms $c\bar{c} \rightarrow \chi_c, J/\Psi$ are explained in [9,10]. Because the color octet contributions depend linearly on the so called color octet matrix elements (COME) which at the moment cannot be calculated but have to be fitted to the data [11,12], the following results should not to be seen as predictions of the corresponding differential cross sections but are required in order to extract the sizes of the COME. The results for the production of the J/Ψ from radiative χ_c -decay in the $k_\perp$ -factorization are shown on the left side of fig. 5 together with experimental data from the Tevatron and a NLO collinear calculation. It is clear from the figure that the $^3S_1^8$ -contribution to χ_c is generally suppressed due to the flat slope of the curve in comparison to the data. The flat curve originates from the additional part of the effective vertex in equation (3) which contains factors $t_{1,2} \approx q_{1,2\perp}^2$ and leads to a strong suppression of the corresponding COME with respect to the results from collinear calculations. Furthermore the Landau-Yang-Theorem, forbidding the production of the $\chi_{cJ=1}$ -state in LO collinear factorization, is not valid in our approach because the gluons are off-shell. This becomes obvious in a rather impressive way on the right hand side of fig. 5 where the individual $\chi_{cJ=1}$ - and $\chi_{cJ=2}$ -contributions are plotted. Fig. 6 shows our results for direct J/Ψ -production in the $k_\perp$ -factorization framework in comparison to data and the NLO color singlet collinear prediction [10]. Although in this case the color octet contributions $^3P_J^8, ^1S_0^8$ play an important role, the $^3S_1^8$ -part is suppressed following the same reasons as in the χ_c -production. This leads again to a strong reduction of the corresponding COME $\langle 0 | \mathcal{O}^{J/\Psi}(^3S_1^8) | 0 \rangle$ with respect to the collinear result. This in turn has a definite influence on the production of polarized J/Ψ [10,13] and could clarify the so far unsolved puzzle concerning the disagreement between recent experimental results and the NLO collinear predictions of transversally polarized charmonia [14–17]. This work was supported by the DFG and the BMBF.




Fig. 6. Direct J/Ψ -production

References

- [1] S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. B242(1990) 97; J.C. Collins and R.K. Ellis, Nucl. Phys. B360(1991) 3
- [2] Ph. Hägler, R. Kirschner, A. Schäfer, L. Szymanowski, O.V. Teryaev, Phys.Rev.D 62 (2000) 071502
- [3] V.S. Fadin and L.N. Lipatov, Nucl. Phys. B477(1996) 767
- [4] J. Kwiecinski, A.D. Martin, A.M. Stasto, Phys.Rev. D56(1997) 3991
- [5] B. Abbott et al., Phys.Lett B487(2000) 264
- [6] F. Abe et al., Phys.Rev. D55(1997) 2546
- [7] Abe et al., Phys.Rev.Lett.79(1997) 578
- [8] Abe et al., CDF Collaboration, FERMILAB-Conf-95/226-E
- [9] Ph. Hägler, R. Kirschner, A. Schäfer, L. Szymanowski, O.V. Teryaev, Phys.Rev.Lett. 86 (2001) 1446
- [10] Ph. Hägler, R. Kirschner, A. Schäfer, L. Szymanowski, O.V. Teryaev, Phys.Rev.D63 (2001) 077501
- [11] P. Cho, A.K. Leibovich, Phys.Rev. D53(1996) 150
- [12] P. Cho, A.K. Leibovich, Phys.Rev. D53(1996) 6203
- [13] F. Yuan, K.-T. Chao, Phys.Rev.Lett. 86(2001) 022002
- [14] M. Beneke, M. Kramer, Phys.Rev. D55(1997) 5269
- [15] E. Braaten, B.A. Kniehl, J. Lee, Phys.Rev. D62(2000) 094005
- [16] T. Affolder et al., Phys.Rev.Lett. 85(2000) 2886
- [17] B.A. Kniehl, J. Lee, Phys.Rev. D62(2000) 114027